



Australian
Centre for
Disease
Control

Australian Respiratory Surveillance Report

Key messages

This report presents a national update on acute respiratory infections, including coronavirus disease 2019 (COVID-19), influenza and respiratory syncytial virus (RSV). It focuses on the current reporting period (6 April to 19 April 2026) and earlier severity reporting periods (up to 5 April 2026). These key messages do not provide detailed information on age distribution, jurisdictional patterns, or comparisons with previous years. This analysis and supporting data are available in the full report.

In the community: In the last fortnight, self-reported influenza-like illness remained relatively stable among people contacting the national health helpline and among those who took part in community surveys. In the last fortnight, COVID-19 and influenza notifications decreased, while RSV notifications increased slightly, consistent with expected seasonal patterns. The continued decrease in COVID-19 and influenza cases at a time where increasing cases might have been expected, and the slowing of the increase in RSV cases observed in the last fortnight may reflect reduced population mixing and changes in testing or health seeking behaviours associated with the recent school holiday period.

In general practice: In the last fortnight, influenza-like illness consultation rates at sentinel general practice sites decreased and are consistent with usual interseasonal levels. Rhinovirus / enterovirus was the most commonly detected respiratory pathogen among tested influenza-like illness patients.

In hospitals: In the most recent severity reporting fortnight, admissions to sentinel hospitals with severe acute respiratory infections remained relatively stable, and most admissions in the last severity fortnight were with RSV. Sentinel intensive care admissions with severe acute respiratory infections increased in the most recent severity reporting period, and most admissions in the last severity period were with RSV. In the last fortnight, the average daily number of intensive care patients in droplet or airborne isolation increased slightly.

Deaths: COVID-19 has been the leading cause of acute respiratory infection mortality across the majority of 2020–2025; however, since August 2025 the number of deaths per month involving influenza (both due to and with) has exceeded the number of deaths involving COVID-19. The mortality burden of acute respiratory infections is highest in older adults.

In laboratories: In the last fortnight, test positivity for SARS-CoV-2 and RSV remained relatively stable, while test positivity for influenza decreased slightly. The SARS-CoV-2 variant under monitoring, NB.1.8.1 was the most common circulating SARS-CoV-2 variant.

Vaccine coverage, effectiveness and match: In the last year, 9.3% of adults have received a COVID-19 vaccine. In 2025, influenza vaccine coverage reached 30.7%; however, 2026 data are not yet available. Most influenza isolates (>87%) are a good match for the 2026 southern hemisphere vaccine components. The National RSV Mother and Infant Protection Program continues. To date, 236,884 Abrysvo doses have been administered, and nirsevimab uptake over the last six months was 4.0% in infants aged up to eight months.

Australian Respiratory Surveillance Report

This report was prepared by Emma Bandy, Nga Nguyen, Jenna Hassall and Shweta Singhal on behalf of the Australian Centre for Disease Control (CDC). We thank the staff and participants from the surveillance systems who contribute data for acute respiratory illness surveillance across Australia.

The report presents a national overview of acute respiratory infections in Australia, drawing information from several different surveillance systems. These surveillance systems help us to understand the distribution of acute respiratory illnesses in the community, the severity of infections including which populations might be at risk, and the impact of acute respiratory illnesses on the community and health system in Australia.

Surveillance indicators presented in this report are based on the [Australian National Surveillance Plan for COVID-19, Influenza, and RSV](#). A summary of data considerations for this report are provided below:

- Due to the dynamic nature of the surveillance systems used in this report, surveillance data are considered preliminary and subject to change as updates are received, with the most recent weeks considered particularly incomplete. Data in this report may vary from data reported in other national reports and reports by states and territories.
- Data in this report are presented by date of event (survey, diagnosis, admission or death) and by the International Organization for Standardization (ISO) week date system, with weeks defined as seven-day periods which begin on a Monday and end on a Sunday. The ISO week date system is used to support trends comparisons over time more effectively. The current reporting period includes 6 April to 19 April 2026 and where comparisons to the previous fortnight are made, this includes 23 March to 5 April 2026.
- In Australia, states and territories (the Australian Capital Territory [ACT], New South Wales [NSW], the Northern Territory [NT], Queensland [Qld], South Australia [SA], Tasmania [Tas], Victoria [Vic] and Western Australia [WA]) report notified cases to the National Notifiable Diseases Surveillance System (NNDSS) based on the [Australian national surveillance case definitions](#). NNDSS data are analysed and reported based on diagnosis date, which is the true onset date of a case if known, otherwise it is the earliest of the specimen date, the notification date or the notification received date. The NNDSS data for this report were extracted on 22 April 2026.
- Notification rates per 100,000 population presented in this report are for the given time period, with population data are based on the Australian Bureau of Statistics (ABS) [Estimated Resident Population \(ERP\) for the reference period June 2024, released 12 December 2024](#) unless stated otherwise.
- To account for the lag in collection and provision of severity data from some surveillance systems, and for the time delay between illness onset and the development of severe disease outcomes, cases with an admission date or a diagnosis date in the last two weeks are excluded from severity analyses for hospitalisations and intensive care admissions. As such, the severity reporting periods are two weeks behind the end of the current reporting period. For this report, severity reporting includes data from 23 March to 5 April 2026 unless specified otherwise. Where comparisons to the previous severity fortnight are made this includes 9 March to 22 March 2026.
- Death registrations from the ABS Provisional Mortality Statistics are now used as the primary data source for measuring acute respiratory infection associated deaths. The ABS mortality data is sourced from the Registry of Births, Deaths and Marriages and is separate from the NNDSS. Registration-based mortality data needs time to be received and processed, and so mortality statistics in this report may lag by at least two months.
- The responsibility for the interpretation and use of the material lies with the reader. The Australian CDC does not accept liability for any injury or loss or damage arising from the use of, or reliance upon, the content of the report. Analysis and reporting outputs were produced using R Statistical Software v4.3.1.
- For further information about this report, including data sources and considerations refer to the [Technical Supplement](#) or contact respiratory.surveillance@cdc.gov.au.


Community surveillance

Community surveillance monitors respiratory illnesses in the community, providing information on the number of people reporting respiratory symptoms, testing practices, and the impact of respiratory illnesses.

Community surveillance includes notification data obtained from laboratory tests for infections. Infections that are diagnosed and notified are only a subset of the total number of infections occurring in the community.

- In the last fortnight (6 April to 19 April 2026), the rate of Healthdirect helpline callers with influenza-like illness (332 per 1,000 callers per fortnight) remained relatively stable compared with the previous fortnight (330 per 1,000 callers per fortnight) (Figure 1).
- Rates of influenza-like illness among helpline callers are similar to the same period in 2025 and 2024 (Figure 1).

Figure 1: Rate of influenza-like illness per 1,000 helpline callers by year and week of call*, Australia†, 22 March 2023 to 19 April 2026




Source: Healthdirect Australia

* Healthdirect data prior to 22 March 2023 are unavailable as prior to this date a different data collection method was used.

† The Healthdirect helpline operates in all states and territories except Qld; therefore influenza-like illness trends will not be representative of Qld and may be underrepresented. See the [Technical Supplement](#) for more information.

- In the last fortnight, the rate of Healthdirect helpline callers with influenza-like illness who were referred to seek urgent medical care (172 per 1,000 callers per fortnight) was relatively similar to the previous fortnight (176 per 1,000 callers per fortnight) (Figure 2).
 - Callers referred to seek urgent medical care include those referred to call triple zero, attend a hospital emergency department, contact a virtual emergency department, urgent care clinic or see a general practitioner within two hours.
- In the last fortnight, referral pathways for influenza-like illness varied across Australia. NSW and Vic had the highest proportion of callers referred to see a general practitioner (GP) or seek other urgent medical care (Figure 2). By comparison, the NT, Tas and WA had a higher proportion of callers who were recommended to attend a hospital emergency department or call triple zero. While the ACT and SA had similar proportions of callers referred to see a GP or seek other urgent medical care and recommended to attend a hospital emergency department or call triple zero (Figure 2).

Figure 2: Proportion of referral categories* for helpline callers with influenza-like illness by jurisdiction† and call date, Australia, 1 January to 19 April 2026




Source: Healthdirect Australia

* See other health professionals category includes pharmacist, dentist, mental health provider, primary maternity care, poison information centre or other.

† The Healthdirect helpline operates in all states and territories except Qld; therefore influenza-like illness referral trends are not provided for Qld. See the [Technical Supplement](#) for more information.

- In the last fortnight, a slightly lower percentage of FluTracking participants reported new fever and cough symptoms (0.9%), than in the previous fortnight (1.1%) (Figure 3). This decline may reflect reduced population mixing and changes in testing or health-seeking behaviour associated with the recent school holiday period.
- The weekly percentage of FluTracking participants reporting new fever and cough symptoms has remained relatively stable across March and April but decreased over the last fortnight. The weekly percentage of FluTracking participants reporting new fever and cough symptoms over the last fortnight was lower than the trend observed in previous years and the historical 5-year average (Figure 3).
- In the last fortnight, the percentage of First Nations FluTracking participants reported new fever and cough symptoms (0.9%) remained the same as the previous fortnight (0.9%). These findings could be impacted by smaller sample sizes and the representativeness of the data. For more detailed trends, please refer to figure 2 in the [FluTracking reports](#).

Figure 3: Age standardised percentage of survey participants reporting new fever and cough symptoms compared with the five-year average* by year and week of survey, Australia, 2023 to 19 April 2026



Source: FluTracking

* From 2020, FluTracking expanded their data capture period to year-round. Data before May and after October for any year before 2020 are not available for historical comparisons. The years 2020 and 2021 are excluded when comparing the current season to historical periods when influenza virus has circulated without public health restrictions. As such, the five-year average includes the years 2019 and 2022 to 2025.

- The average percentages of FluTracking participants reporting taking three or more days off work or normal duties, or seeking medical advice or care, due to fever and cough symptoms in 2026 are both lower than the same period in previous years (Table 1).

Table 1: Percentage of FluTracking participants reporting new fever and cough symptoms plus three or more days off work or normal duties or seeking medical advice or care*, Australia, up to 5 April† for 2023–2026

	2023	2024	2025	2026
Reported three or more days off work or normal duties	50.6%	51.0%	47.9%	44.5%
Reported seeking medical advice or care*	35.6%	32.1%	32.0%	30.2%


Source: FluTracking

* Includes those who sought medical advice from a general practitioner, Aboriginal and Torres Strait Islander health clinic, COVID-19 clinic, emergency department, or were admitted to hospital for fever and cough.

† While FluTracking data are collected in real time, data presented here are subject to a two week reporting delay to account for the time delay between illness onset and the development of severe disease outcomes.


- Self-reported severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) polymerase chain reaction (PCR) positivity has varied considerably, peaking in late February at 27.6% per week (Figure 4a). Self-reported SARS-CoV-2 rapid antigen test (RAT) positivity has remained relatively low and stable, with some week-on-week variation observed (Figure 4b).
- Self-reported influenza PCR positivity declined from 40.0% per week in late January to 5.9% in mid-April with some week-on-week variation observed throughout 2026 (Figure 4a). Self-reported influenza RAT positivity has followed a similar decline from January down to 3.6% per week in late March and has since remained relatively low and stable (Figure 4b).
- Self-reported RSV PCR positivity has fluctuated in 2026, peaking at 11.8% per week in mid-April (Figure 4a). Self-reported RSV RAT positivity has also fluctuated, peaking at 11.8% per week in late January with a second, smaller peak of 10.2% per week in mid-February, but has since remained relatively low and stable (Figure 4b).
- For more detailed testing and self-reported positivity trends, please refer to the [FluTracking reports](#).

Figure 4a: Self-reported PCR test positivity* among FluTracking participants with fever and cough symptoms by pathogen week of survey, Australia, 1 January to 19 April 2026



Source: FluTracking
 * Denominator is based on participants who self-reported fever and cough symptoms and had a PCR test. Please refer to the [Technical Supplement](#) for more details.

Figure 4b: Self-reported RAT positivity* among FluTracking participants with fever and cough symptoms by pathogen and week of survey, Australia, 1 January to 19 April 2026



Source: FluTracking
 * Denominator is based on participants who self-reported fever and cough symptoms and had a RAT. Please refer to the [Technical Supplement](#) for more details.

- In the last fortnight (6 April to 19 April 2026), there was a 17.0% decrease in COVID-19 cases, a 18.5% decrease in influenza cases, and a 3.8% increase in RSV cases, compared with the previous fortnight. The continued decrease in COVID-19 and influenza cases at a time where increasing cases might have been expected, and the slowing of the increase in RSV cases observed in the last fortnight may reflect reduced population mixing and changes in testing or health seeking behaviour associated with the recent school holiday period.

Table 2: Notified cases and notification rate per 100,000 population by disease, five-year age group, and jurisdiction*, Australia, 1 January to 19 April 2026


Age group (years)	COVID-19			Influenza			RSV		
	Reporting period (n)	Year to date (n)	Year to date (rate)	Reporting period (n)	Year to date (n)	Year to date (rate)	Reporting period (n)	Year to date (n)	Year to date (rate)
0–4	673	4,898	325	264	3,357	222	2,723	13,081	867
5–9	132	1,675	104	178	2,689	167	282	1,390	86
10–14	58	1,367	82	96	1,980	118	93	662	40
15–19	92	1,048	63	74	1,963	118	103	707	43
20–24	129	991	55	100	1,927	108	120	683	38
25–29	160	1,147	57	85	1,583	79	119	752	38
30–34	211	1,513	74	93	1,410	69	180	879	43
35–39	215	1,573	79	99	1,358	68	167	818	41
40–44	194	1,450	78	98	1,351	73	137	708	38
45–49	159	1,181	73	94	1,152	71	134	797	49
50–54	156	1,165	69	83	1,092	65	171	944	56
55–59	144	1,132	74	88	1,147	75	191	1,112	73
60–64	173	1,106	72	79	1,170	76	233	1,123	73
65–69	152	1,061	78	86	1,203	88	238	1,110	82
70–74	183	1,232	105	79	1,151	98	239	1,157	99
75+	782	5,389	249	268	3,315	153	798	3,659	169
Jurisdiction									
ACT	51	286	60	14	310	65	39	188	40
NSW	1,803	12,344	145	736	9,695	114	3,325	14,314	169
NT	9	141	55	20	542	212	54	636	249
Qld	753	7,304	131	535	8,675	155	1,434	8,836	158
SA	312	2,318	123	113	1,623	86	112	819	44
Tas	49	324	56	10	303	53	39	298	52
Vic	548	4,298	62	352	4,976	71	700	3,290	47
WA	89	924	31	84	1,737	59	225	1,204	41
Total	3,614	27,939	103	1,864	27,861	102	5,928	29,585	109

Source: National Notifiable Diseases Surveillance System (NNDSS). RSV notification data are unavailable for Tas from 17 April 2026.

* Total includes cases with missing age.


- In the last fortnight, there were 3,614 COVID-19 cases, a 17.0% decrease from 4,352 cases notified in the previous fortnight (Table 2; Figure 5).
- In the year to date, there have been 27,939 COVID-19 cases, 47.8% fewer than the 53,534 cases notified over the same period in 2025 (Table 2; Figure 5). There is still no consistent seasonal pattern for COVID-19.
- In the last fortnight, COVID-19 notification rates varied across jurisdictions compared with the previous fortnight, with COVID-19 notification rates decreasing in NSW, Qld, SA and WA but increasing in the ACT and Tas (Figure 6).

Figure 5: Notified COVID-19 cases by year and week of diagnosis, Australia, 2023 to 19 April 2026



Source: National Notifiable Diseases Surveillance System (NNDSS)


Figure 6: Notification rates per 100,000 population for COVID-19 cases by state or territory and week of diagnosis, Australia, 1 January to 19 April 2026



Source: National Notifiable Diseases Surveillance System (NNDSS)

- In the last fortnight, there were 1,864 influenza cases, a 18.5% decrease from 2,286 cases notified in the previous fortnight (Table 2; Figure 7).
- In the year to date, there have been 27,861 influenza cases, 56.6% fewer than the 64,238 cases notified over the same period in 2025 (Table 2; Figure 7). This decrease could potentially reflect greater population immunity following the higher than usual influenza activity in late 2025 and reduced overseas importations due to earlier influenza activity in some northern hemisphere countries in their 2025-26 influenza season.
- In the last fortnight, influenza notification rates decreased or remained relatively stable across most jurisdictions compared with the previous fortnight, except in Vic where notification rates increased slightly (Figure 8).


Figure 7: Notified influenza cases and five-year average* by year and week of diagnosis, Australia, 2023 to 19 April 2026



Source: National Notifiable Diseases Surveillance System (NNDSS)

* The years 2020 and 2021 are excluded when comparing the current season to historical periods when influenza virus has circulated without public health restrictions. As such, the five-year average includes the years 2019 and 2022 to 2025. Please refer to the [Technical Supplement](#) for interpretation of the five-year average.


Figure 8: Notification rates per 100,000 population for influenza cases by state or territory and week of diagnosis, Australia, 1 January to 19 April 2026



Source: National Notifiable Diseases Surveillance System (NNDSS)

- In the last fortnight, there were 1,500 influenza A cases, a 17.5% decrease from 1,819 influenza A cases notified in the previous fortnight, and there were 348 influenza B cases, a 23.0% decrease from 452 influenza B cases notified in the previous fortnight.
- Among influenza A cases with subtype information available there were:
 - 13 influenza A(H1N1) cases notified in the last fortnight, a 31.6% decrease from 19 cases notified in the previous fortnight.
 - 42 influenza A(H3N2) cases notified in the last fortnight, a 53.3% decrease from 90 cases notified in the previous fortnight.
- In the year to date, influenza A(Unsubtyped) has accounted for most cases across all age groups, with the number of influenza B notifications remaining relatively consistent since early February (Figure 9). Trends in influenza subtypes are influenced by differences in the number and selection of influenza samples that undergo typing across age groups and healthcare settings.


Figure 9: Notified influenza cases by influenza subtype, age group*, and week of diagnosis, Australia, 1 January to 19 April 2026



Source: National Notifiable Diseases Surveillance System (NNDSS)
 * Axis varies between age groups.

- In the year to date, influenza A(Unsubtyped) has accounted for most influenza cases across all jurisdictions; however, an increasing number of influenza B notifications has been observed in most jurisdictions since early February, most notably in the NT (Figure 10).
- Trends in influenza subtypes are influenced by jurisdictional differences in the number and selection of influenza samples that undergo typing.

Figure 10: Notified influenza cases by influenza subtype, jurisdiction*, and week of diagnosis, Australia, 1 January to 19 April 2026




Source: National Notifiable Diseases Surveillance System (NNDSS)

* Axis varies between jurisdictions.


- In the last fortnight, there were 5,928 RSV cases, a 3.8% increase from 5,711 cases notified in the previous fortnight. RSV cases have been steadily increasing since late January, indicating that the 2026 RSV season has commenced (Table 2; Figure 11).
- In the year to date, there have been 29,585 RSV cases, 15.5% fewer than the 34,996 cases notified over the same period in 2025 (Table 2; Figure 11).
- In the last fortnight, RSV notification rates varied across jurisdictions compared with the previous fortnight, with RSV notification rates decreasing in the NT, Qld and SA and but increasing in NSW, Vic and WA (Figure 12). RSV notification data were unavailable for Tas from 17 April 2026.

Figure 11: Notified RSV cases by year and week of diagnosis, Australia, 2023 to 19 April 2026



Source: National Notifiable Diseases Surveillance System (NNDSS)

Figure 12: Notification rates per 100,000 population for RSV cases by state or territory and week of diagnosis, Australia, 1 January to 19 April 2026




Source: National Notifiable Diseases Surveillance System (NNDSS). RSV notification data are unavailable for Tas from 17 April 2026.

Primary care surveillance

Primary care surveillance monitors the number and characteristics of people who have presented to a general practice with influenza-like illness and provides insight on the different respiratory pathogens that are causing illness in the community.

- In the last fortnight (6 April to 19 April 2026), there were fewer general practice consultations for influenza-like illness (3.1 notifications per 1,000 consultations per fortnight) compared to the previous fortnight (3.8 notifications per 1,000 consultations per fortnight) (Figure 13).
- From late January, influenza-like illness consultation rates gradually increased, before gradually declining since late March. In the last fortnight, influenza-like illness consultation rates have generally been lower than the trend observed in previous years and the historical 5-year average (Figure 13).

Figure 13: Rate of influenza-like illness notifications per 1,000 consultations per week in sentinel general practice sites compared with the five-year average by year and week of consultation*†, Australia, 2023 to 19 April 2026




Source: Australian Sentinel Practices Research Network (ASPREN)

* The years 2020 and 2021 are excluded when comparing the current season to historical periods when influenza virus has circulated without public health restrictions. As such, the five-year average includes the years 2019 and 2022 to 2025. Please refer to the [Technical Supplement](#) for interpretation of the five-year average.

† Please refer to the [Technical Supplement](#) for notes on impact of COVID-19 on ASPREN data.

- In the last fortnight, 54.4% (43/79) of people attending general practice with influenza-like illness who were tested have then tested positive for a respiratory pathogen.
- In the last fortnight, rhinovirus / enterovirus (48.8%; 21/43) was the most commonly detected pathogen, followed by parainfluenza (18.6%; 8/43) and influenza (11.6%; 5/43) (Figure 14).
- In the year to date, 57.5% (307/534) of people attending general practice with influenza-like illness who were tested have then tested positive for a respiratory pathogen.
- In the year to date, rhinovirus / enterovirus (51.8%; 159/307) has been the most commonly detected pathogen, followed by influenza (12.1%; 37/307), parainfluenza (11.4%; 35/307), SARS-CoV-2 (6.8%; 21/307), RSV (6.5%; 20/307) (Figure 14).

Figure 14: Number of samples tested for respiratory pathogens among people with influenza-like illness attending sentinel general practice sites by respiratory pathogen and week of specimen collection, Australia, 1 January to 19 April 2026



Source: Australian Sentinel Practices Research Network (ASPREN)


Note: All ASPREN swab samples are transported to the SA Pathology laboratory in Adelaide to be tested for viral and bacterial respiratory pathogens via a multiplex real-time reverse transcription polymerase chain reaction (RT-PCR) assay using in-house primers.

Hospital-based surveillance

Hospital-based surveillance monitors persons with more severe illness who have been admitted to hospital for their respiratory illness (severe acute respiratory infections). Hospital-based surveillance also measures the ability of the health system to cope with the number of severe acute respiratory infection admissions to ensure delivery of safe, timely and quality health care.

- In the last severity reporting period (23 March to 5 April 2026), a similar number of patients were admitted to a sentinel hospital with a severe acute respiratory infection (n=220), compared with the previous severity reporting period (n=224).
 - In the last severity reporting period, at sentinel hospitals there were 11.1% fewer admissions with COVID-19 (from 72 to 64), 60.6% fewer admissions with influenza (from 33 to 13), and 20.2% more admissions with RSV (from 119 to 143), compared to the previous severity reporting period.
- In the year to date for severity reporting (1 January to 5 April 2026), there have been 1,394 admissions with severe acute respiratory infections at sentinel hospitals. Most patients with a severe acute respiratory infection have been admitted with RSV (n=579) followed by influenza (n=418) (Figure 15).


Figure 15: Total number of patients (children and adults) admitted with a severe acute respiratory infection to sentinel hospitals by disease and week of admission, Australia, 1 January to 5 April 2026



Source: Influenza Complications Alert Network (FluCAN)

- Patients admitted to sentinel hospitals with influenza have mostly been admitted with influenza A (91.6%; 383/418), while 8.1% (34/418) were admitted with influenza B.
 - Most hospital admissions with influenza A have been with influenza A(Unsubtyped) (75.5%; 289/383), followed by influenza A(H3N2) (19.1%; 73/383) and then influenza A(H1N1) (5.5%; 21/383).
- In the year to date for severity reporting, influenza A (Unsubtyped) was the most commonly detected influenza type in all age groups. Influenza B represented a smaller number of admissions overall, with a relatively higher contribution observed in younger age groups compared with adults (Figure 16).

Figure 16: Number of patients admitted with influenza to sentinel hospitals by influenza subtype, age group*, and week of admission, Australia, 1 January to 5 April 2026



Source: Influenza Complications Alert Network (FluCAN)

* Axis varies between age groups. The age distribution of admissions with influenza may not reflect the age distribution of all patients.

- More children (those aged 16 years and younger) have been admitted with RSV to sentinel hospitals than with COVID-19 or influenza (Table 3a).
- Children admitted to sentinel hospitals with influenza tended to be older than children admitted with COVID-19 or RSV (Table 3a).
- Children admitted to sentinel hospitals with RSV, COVID-19 or influenza had similar lengths of hospital stays (Table 3a).
- Sadly, a small number of children admitted to sentinel hospitals with influenza or RSV have died (Table 3a).

Table 3a: Demographic characteristics and outcomes for children admitted with a severe acute respiratory infection to a sentinel hospital by disease*†‡, Australia, 1 January to 5 April 2026

	COVID-19 Year to date for severity reporting (n=197)	Influenza Year to date for severity reporting (n=221)	RSV Year to date for severity reporting (n=467)
Age (years)			
Median [IQR]	1 [0-5]	3 [1-7]	1 [0-2]
Age group (years)			
< 6 months	47 (23.9%)	27 (12.2%)	87 (18.6%)
6 months – 4 years	98 (49.7%)	108 (48.9%)	347 (74.3%)
5–16 years	52 (26.4%)	86 (38.9%)	33 (7.1%)
Indigenous status			
Aboriginal and Torres Strait Islander	13 (6.6%)	25 (11.3%)	65 (13.9%)
Length of hospital stay (days)†			
Median [IQR]	2 [1-2]	1 [1-2]	2 [1-3]
Patient admission location‡			
Admitted to hospital ward	192 (97.5%)	211 (95.5%)	444 (95.1%)
Admitted to intensive care directly	5 (2.5%)	10 (4.5%)	23 (4.9%)
Discharge status†			
Alive	172 (87.3%)	210 (95.0%)	401 (85.9%)
Died	-	1 (0.5%)	1 (0.2%)
Incomplete/missing	25 (12.7%)	10 (4.5%)	65 (13.9%)

Source: Influenza Complications Alert Network (FluCAN)

* Does not include patients with missing age; therefore, the sum of age-specific totals above may not equal the total number of patients.

† For patients who are still in hospital data may not be complete; therefore, these data are not included in the length of stay or discharge status. In addition, length of stay data excludes patients that acquired their infection in hospital.

‡ Admission location reflects the initial admission ward. Some patients may be initially admitted to general ward then later admitted to an intensive care and this is not reflected here. Does not include patients with missing admission location; therefore, the sum of admission location specific totals above may not equal the total number of patients.

The Paediatric Active Enhanced Disease Surveillance (PAEDS) network carries out enhanced sentinel hospital surveillance for some acute respiratory infections or conditions in children. PAEDS data for acute respiratory infections in children are presented in the Australian Respiratory Surveillance Reports in the sentinel hospital data from FluCAN. For additional information on COVID-19, influenza or RSV in children please visit the [PAEDS](#) webpages and dashboards.

- More adults (those aged 17 years and over) have been admitted with influenza or COVID-19 to sentinel hospitals than with RSV (Table 3b).
- A greater proportion of adults aged 65 years and over have been admitted to sentinel hospitals with COVID-19, influenza and RSV compared to adults aged 17–64 years (Table 3b).
- Adults admitted to sentinel hospitals with COVID-19, influenza or RSV had similar lengths of hospital stays (Table 3b).
- Sadly, a small number of adults admitted to sentinel hospitals with severe acute respiratory infections have died (Table 3b).

Table 3b: Demographic characteristics and outcomes for adults admitted with a severe acute respiratory infection to a sentinel hospital by disease*†‡, Australia, 1 January to 5 April 2026

	COVID-19 Year to date for severity reporting (n=200)	Influenza Year to date for severity reporting (n=197)	RSV Year to date for severity reporting (n=112)
Age (years)			
Median [IQR]	72 [59-82]	68 [50-80]	69 [56-80]
Age group (years)			
17–64 years	72 (36.0%)	81 (41.1%)	40 (35.7%)
65 years and over	128 (64.0%)	116 (58.9%)	72 (64.3%)
Indigenous status			
Aboriginal and Torres Strait Islander	21 (10.5%)	17 (8.6%)	13 (11.6%)
Length of hospital stay (days)†			
Median [IQR]	4 [3-8]	3 [1-6]	4 [2-7]
Patient admission location‡			
Admitted to hospital ward	192 (96.0%)	180 (91.4%)	106 (94.6%)
Admitted to intensive care directly	8 (4.0%)	17 (8.6%)	6 (5.4%)
Discharge status†			
Alive	137 (68.5%)	168 (85.3%)	86 (76.8%)
Died	7 (3.5%)	4 (2.0%)	5 (4.5%)
Incomplete/missing	56 (28.0%)	25 (12.7%)	21 (18.8%)

Source: Influenza Complications Alert Network (FluCAN)


* Does not include patients with missing age; therefore, the sum of age-specific totals above may not equal the total number of patients.

† For patients who are still in hospital data may not be complete; therefore, these data are not included in the length of stay or discharge status. In addition, length of stay data excludes patients that acquired their infection in hospital.

‡ Admission location reflects the initial admission ward. Some patients may be initially admitted to general ward then later admitted to an intensive care and this is not reflected here. Does not include patients with missing admission location; therefore, the sum of admission location specific totals above may not equal the total number of patients.


- In the last severity reporting period for sentinel intensive care (9 March to 5 April 2026), more patients have been admitted to a sentinel intensive care with a severe acute respiratory infection (n=104), than in the previous severity reporting period (n=83) (Figure 17).
- In the year to date for severity reporting (1 January to 5 April 2026), most patients were admitted to sentinel intensive care with influenza, followed by RSV (Figure 17; Table 4).

Figure 17: Number of patients admitted with severe acute respiratory infections to a sentinel intensive care by disease and week of admission, Australia, 1 January to 5 April 2026



Source: Short Period Incidence Study of Severe Acute Respiratory Infection (SPRINT-SARI) Australia
 Note: The sum of pathogen-specific totals above may not equal the total number of severe acute respiratory infection patients due to co-infections.

Figure 18: Number of patients admitted with severe acute respiratory infections to a sentinel intensive care by disease and age group*, Australia, 1 January to 5 April 2026



Source: Short Period Incidence Study of Severe Acute Respiratory Infection (SPRINT-SARI) Australia.
 Note: The sum of pathogen-specific totals above may not equal the total number of severe acute respiratory infection patients due to co-infections.
 * The age distribution of severe acute respiratory infection intensive care admissions may not reflect the age distribution of all patients.

- In the year to date for severity reporting, most admissions have been among children aged 0–9 years followed by adults aged 50 years and over (Figure 18; Table 4).
- The proportion of admissions requiring invasive mechanical ventilation was higher in patients with influenza and parainfluenza; however, the length of invasive mechanical ventilation was longest in patients with *Mycoplasma pneumoniae* (Table 4). The length of intensive care stay has been relatively

similar across pathogens; however, comparisons across are limited by small numbers for some pathogens.

- Most patients with a severe acute respiratory infection have been discharged home. Sadly, a number of patients have died in hospital (Table 4).

Table 4: Demographic characteristics and outcomes of patients admitted with a severe acute respiratory infection to a sentinel intensive care by disease*†, Australia, 1 January to 5 April 2026

	COVID-19 Year to date for severity reporting (n=69)	hMPV Year to date for severity reporting (n=43)	Influenza Year to date for severity reporting (n=95)	Mycoplasma pneumoniae Year to date for severity reporting (n=7)	Parainfluenza Year to date for severity reporting (n=53)	RSV Year to date for severity reporting (n=70)
Age (years)						
Median [IQR]	65 [42–78]	44 [9–64]	60 [40–73]	55 [46–76]	54 [9–68]	34 [5–61]
Indigenous status						
Aboriginal and Torres Strait Islander	4 (5.8%)	5 (11.6%)	13 (13.7%)	–	6 (11.3%)	13 (18.6%)
Non-Indigenous	65 (94.2%)	38 (88.4%)	82 (86.3%)	7 (100.0%)	47 (88.7%)	57 (81.4%)
Received invasive mechanical ventilation						
Number (%)	19 (27.5%)	8 (18.6%)	31 (32.6%)	2 (28.6%)	18 (34.0%)	14 (20.0%)
Length of invasive mechanical ventilation (days)*						
Median [IQR]	3 [1–8]	7 [3–8]	3 [1–10]	11 [7–15]	2 [1–10]	3 [1–5]
Length of intensive care stay (days)*						
Median [IQR]	3 [2–4]	3 [1–5]	3 [2–7]	2 [1–2]	4 [2–7]	3 [1–4]
Length of hospital stay (days)*						
Median [IQR]	6 [3–14]	7 [4–10]	7 [4–13]	4 [4–5]	8 [4–13]	6 [3–10]
Patient outcome†						
Ongoing care in intensive care	8 (11.6%)	–	1 (1.1%)	1 (14.3%)	3 (5.7%)	3 (4.3%)
Ongoing care in hospital ward	4 (5.8%)	1 (2.3%)	2 (2.1%)	–	2 (3.8%)	4 (5.7%)
Transfer to other hospital / facility	6 (8.7%)	6 (14.0%)	14 (14.7%)	–	5 (9.4%)	5 (7.1%)
Discharged home	41 (59.4%)	32 (74.4%)	65 (68.4%)	6 (85.7%)	35 (66.0%)	53 (75.7%)
Died in hospital	10 (14.5%)	4 (9.3%)	12 (12.6%)	–	8 (15.1%)	5 (7.1%)

Source: Short Period Incidence Study of Severe Acute Respiratory Infection (SPRINT-SARI) Australia


Note: The sum of pathogen-specific totals above may not equal the total number of severe acute respiratory infection patients due to co-infections.

* For patients receiving ongoing care in intensive care data may not be complete; therefore, data are not included in the length of ventilation or stay.

† Patients who have been admitted with no discharge information for less than 90 days have been assumed to have ongoing care in the hospital. Patients who have no outcome entered or have been admitted for more than 90 days with no discharge information have been treated as missing.


- In the last fortnight (6 April to 19 April 2026), there were an average of 173 intensive care patients in droplet or airborne isolation for any suspected or confirmed respiratory pathogen each day, a 1.8% increase from an average of 170 patients in isolation each day reported in the previous fortnight (Figure 19).
 - Suspected or confirmed respiratory pathogens may include nationally notifiable conditions such as COVID-19, influenza, RSV or pertussis (Whooping cough) but also other non-notifiable respiratory pathogens like adenovirus, hMPV, parainfluenza, rhinovirus or bacterial infections causing atypical pneumonias.
- In the last fortnight (6 April to 19 April 2026), there were an average of 82 intensive care staff unavailable to work due to illness each day, a 10.8% increase from an average of 74 staff unavailable each day reported in the previous fortnight (Figure 19).
- In the last fortnight, the average number of intensive care patients in droplet or airborne isolation for any suspected or confirmed respiratory pathogen each day varied across jurisdictions compared with the previous fortnight, with the average number of patients in isolation each day slightly decreasing in SA and Vic but slightly increasing in NSW and Qld (Figure 20).
- In the last fortnight, the average number of intensive care staff unavailable to work due to illness each day remained relatively stable across all jurisdictions compared with the previous fortnight (Figure 20).

Figure 19: Weekly average daily occupancy of intensive care patients in droplet or airborne isolation for any suspected or confirmed respiratory pathogen and the weekly average daily number of intensive care staff unavailable to work due to illness by week of report*, Australia, 1 January to 19 April 2026



Source: Critical Health Resource Information System (CHRIS)
 * Intensive care staff include both medical and nursing staff. Staff unavailability will be underestimated in NSW as most public hospitals in NSW do not report staff unavailability.

Figure 20: Weekly average daily occupancy of intensive care patients in droplet or airborne isolation for any suspected or confirmed respiratory pathogen and the weekly average daily number of intensive care staff unavailable to work due to illness by jurisdiction and week of report^{††}, Australia, 1 January to 19 April 2026



Source: Critical Health Resource Information System (CHRIS)

* Axis varies between jurisdictions.

† NSW isolation data from public hospitals includes all patients occupying intensive care beds in isolation precautions, including those in contact isolation precautions, rather than just droplet or airborne isolation precautions, which will overestimate the average number of patients occupying intensive care beds in droplet or airborne isolation in NSW. For this reason, NSW data may not be comparable to data from other jurisdictions.

‡ Intensive care staff include both medical and nursing staff. Staff unavailability will be underestimated in NSW as most public hospitals in NSW do not report staff unavailability. For this reason, NSW data may not be comparable to data from other jurisdictions.


Mortality surveillance

Death registrations can provide information on the scale and severity of disease associated with acute respiratory infections. An acute respiratory associated death is one where the death was *due to* the disease (the illness has caused terminal complications such as pneumonia) or the person has died *with* the disease (a person has died from another cause but the illness still contributed significantly to death). For more information refer to the [Technical Supplement](#).

There has not been an update to the Provisional Mortality Statistics. The mortality surveillance data presented here have not been updated since the previous report.

- COVID-19 has been the leading cause of acute respiratory infection related mortality across the majority of 2020–2025; however, since August 2025 the number of deaths per month involving influenza (both *due to* and *with*) has exceeded COVID-19 deaths. Deaths involving influenza remain unseasonably high, with 133 deaths in December 2025 and 101 deaths in January 2026.
- Deaths involving COVID-19 (both *due to* and *with*) tend to peak twice a year - between November and January and then again between May and August. Preliminary data does not show a 2025–2026 summer peak, despite a small increase in deaths involving COVID-19 in January 2026.
- In 2025 influenza mortality rates were higher for both Aboriginal and Torres Strait Islander and non-Indigenous people than in 2022–2024. In contrast, in 2025 COVID-19 mortality rates among Aboriginal and Torres Strait Islander people were four times lower than in 2024.
- These acute respiratory infections are more likely to cause death in older age groups.

Figure 21a: Provisional numbers of deaths *due to* an acute respiratory infection*† by month, year, and disease, Australia, 1 January 2024 to 28 February 2026




Source: Australian Bureau of Statistics (ABS), [Deaths due to acute respiratory infections in Australia](#), released 30 March 2026.

* Axis varies between acute respiratory infections.

† Data is provisional and subject to change. It can take several weeks for death registrations to be reported, processed, coded, validated, and tabulated. Therefore, the data shown here may be incomplete. Data for some months were not published by the ABS due to small counts, and therefore not reported here. Data includes all deaths (both doctor and coroner certified) that occurred and were registered by 28 February 2026.

- Deaths *due to* COVID-19 increased slightly in January 2026 but remains at very low levels compared to previous years. The 1,715 deaths *due to* COVID-19 in 2025 are well below both 2024 (3,905 deaths) and 2023 (4,611 deaths) (Figure 21a).
- Deaths *due to* influenza fell in January 2026 but remain high for the time of year. There were 1,449 deaths *due to* influenza in 2025, above the 1,276 deaths in 2017 and the 1,072 deaths in 2019, which were recent high mortality years for influenza (Figure 21a). While influenza deaths were high in 2025, this is expected when there are higher case numbers, and other surveillance systems have not indicated that disease severity was greater than in previous years.
- Deaths *due to* RSV remained at low levels in January 2026, similar to the number of deaths in November and December 2025 (Figure 21a).
- Deaths *with* COVID-19 were slightly higher in January 2026 than December 2025, but remain at low levels compared to previous years (Figure 21b).
- Deaths *with* influenza or *with* RSV decreased in January 2026 and were similar to the number of deaths *with* COVID-19 (Figure 21b).
- The number of deaths *with* RSV was higher in 2025 (393 deaths) than in 2024 (341 deaths) and in 2023 (279 deaths) (Figure 21b).

Figure 21b: Provisional numbers of deaths *with* an acute respiratory infection*† by month, year, and disease, Australia, 1 January 2024 to 28 February 2026



Source: Australian Bureau of Statistics (ABS), [Deaths due to acute respiratory infections in Australia](#), released 30 March 2026.

* Axis varies between acute respiratory infections.


† Data is provisional and subject to change. It can take several weeks for death registrations to be reported, processed, coded, validated, and tabulated. Therefore, the data shown here may be incomplete. Data for some months were not published by the ABS due to small counts, and therefore not reported here. Data includes all deaths (both doctor and coroner certified) that occurred and were registered by 28 February 2026.

Laboratory surveillance

Sentinel laboratory surveillance monitors the percentage of tests with the notifiable condition detected (i.e. test positivity). It also provides information on what pathogens are circulating, potential changes in the pathogens that might affect their infectiousness, severity, ability to evade vaccine and/or infection-acquired immunity, or resistance to antivirals.

- In the last fortnight (6 April to 19 April 2026), the percentage of SARS-CoV-2 tests that were positive remained stable (from 2.5% to 2.5%), the percentage of influenza tests that were positive decreased (from 1.4% to 1.0%) and the percentage of RSV tests that were positive remained stable (from 1.3% to 1.3%) (Figure 22).

Figure 22: Number of tests positive (bars) and percentage of tests positive (line) for SARS-CoV-2, influenza or RSV of those specimens tested by sentinel laboratories by week of report*†, Australia, 1 January to 19 April 2026




Source: Sentinel laboratories, including National Influenza Centres

* Number of specimens tested excludes data from WA as testing denominator data are different for the three pathogens in Western Australia.

† A small minority of total samples from Victoria are tested only by respiratory panel (influenza, parainfluenza, adenovirus, human metapneumovirus, seasonal coronaviruses, RSV, and some picornaviruses) but not for SARS-CoV-2. These minority samples include only forensic materials; all other samples are tested by respiratory panel and SARS-CoV-2 assay.

- There were 86 SARS-CoV-2 sequences uploaded to AusTrakka with dates of collection in the last 28 days (23 March to 19 April 2026). These sequences were from NSW, SA and Tas with the most recent date of collection from 6 April 2026. The low number of sequences uploaded to AusTrakka in the last 28 days is likely due to the reduced number of COVID-19 cases and changes in sequencing capacity and priorities, due to issues with very limited number of sequences and representativeness of the sequences.
- Most sequences were assigned to the BA.2.86 sub-lineage within B.1.1.529 (Omicron) or recombinants consisting of one or more Omicron sub-lineages (Figure 23a/b). In the last 28 days:
 - 58.1% (50/86) of sequences were recombinant or recombinant sub-lineages, the most common including NB.1.8.1 (n=29) and XFG (n=21)
 - 31.4% (27/86) of sequences were identified as BA.3, specifically BA.3.2
 - 7.0% (6/86) of sequences were from the sub-sub-lineages JN.1 (BA.2.86.1.1), specifically KP.3.1.1
 - there were no BA.1, BA.4, BA.5 or other BA.2 sub-sub-lineage sequences.
- NB.1.8.1 was the most common sub-lineage in the last 28 days, accounting for 33.7% (29/86) of sequences (Figure 23a).
- The World Health Organization (WHO) have identified certain sub-sub-lineages and recombinants as variants under monitoring (VUM) because of their epidemiological, pathological, or immunological features of concern. A select number are highlighted below due to their relevance in the Australian context. There are:
 - 4,509 KP.3.1.1 sequences in AusTrakka, with 6 identified in the last 28 days
 - 3,449 NB.1.8.1 sequences in AusTrakka, with 29 identified in the last 28 days
 - 1,157 XFG sequences in AusTrakka, with 21 identified in the last 28 days
 - 499 BA.3.2 sequences in AusTrakka, with 27 identified in the last 28 days

Figure 23a: SARS-CoV-2 Omicron sub-lineage* sequences by sample collection date, showing proportions of sequences per week^{†‡}, Australia, 1 January to 19 April 2026




Source: AusTrakka

* Some sub-sublineages are shown alongside their parent lineage, but not included in the parent lineage totals. For instance, KP.2 and KP.3 are sub-sub lineages of JN.1, so the total of JN.1 sequences will be higher than shown in the corresponding colour alone, and should include the KP.2 and KP.3 totals.

† Sequences in AusTrakka aggregated by week and reported based on date of sample collection, not date of sequencing.

‡ Proportions in Figure 23a may not be representative when sequence numbers are small; refer to Figure 23b. Data for earlier weeks may change between reporting periods as sequences with older collection dates are uploaded. These numbers are not equivalent to number of cases, as there are many cases which may not be sequenced. Non-VOI and non-VUM Omicron sub-lineages have been collapsed into parent lineages BA.1, BA.2, BA.3, BA.4 and BA.5.

Figure 23b: SARS-CoV-2 Omicron sub-lineage* sequences by sample collection date, showing the count of sequences per week^{†‡}, Australia, 1 January 2022 to 19 April 2026



Source: AusTrakka

* Some sub-sublineages are shown alongside their parent lineage, but not included in the parent lineage totals. For instance, KP.2 and KP.3 are sub-sub lineages of JN.1, so the total of JN.1 sequences will be higher than shown in the corresponding colour alone, and should include the KP.2 and KP.3 totals.

† Sequences in AusTrakka aggregated by week and reported based on date of sample collection, not date of sequencing.

‡ Data for earlier weeks may change between reporting periods as sequences with older collection dates are uploaded. These numbers are not equivalent to number of cases, as there are many cases which may not be sequenced. Non-VOI and non-VUM Omicron sub-lineages have been collapsed into parent lineages BA.1, BA.2, BA.3, BA.4 and BA.5.

- In the year to date, the WHO Collaborating Centre for Reference and Research on Influenza has antigenically characterised 341 influenza viruses from Australia (Table 5), of which:
 - 9.4% (32/341) have been influenza A(H1N1)
 - 80.1% (273/341) have been influenza A(H3N2)
 - 10.6% (36/341) have been influenza B/Victoria.
- In the year to date, there have been no influenza B/Yamagata viruses characterised (Table 5). The last influenza B/Yamagata virus characterised in Australia was in a sample from 2020.
- In the year to date none of the samples tested demonstrated highly reduced inhibition to oseltamivir or zanamivir.

Table 5: Australian influenza viruses typed by haemagglutination inhibition assay and jurisdiction*†, 1 January to 19 April 2026

Strain	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total
A(H1N1)	2	8	3	3	1	6	7	2	32
A(H3N2)	21	31	74	12	13	24	89	9	273
B/Victoria lineage	2	7	19	1	1	0	5	1	36
B/Yamagata lineage	0	0	0	0	0	0	0	0	0
Total	25	46	96	16	15	30	101	12	341

Source: World Health Organization (WHO) Collaborating Centre for Reference and Research on Influenza

*Viruses tested by the WHO Collaborating Centre for Reference and Research on Influenza are not necessarily a random sample of all those in the community and early-year data may be based on limited samples received. There may be up to a month delay on reporting of samples.

† Jurisdiction indicates the residential location for the individual tested, not the submitting laboratory.

Vaccine coverage, effectiveness and match

Vaccine coverage, effectiveness and match for acute respiratory infections are monitored from several data sources in Australia. Refer to the [Technical Supplement](#) for more information.

Vaccine coverage

- Nationally, 3.4% of adults (aged 18 years and over) have received a COVID-19 vaccine in the last six months (Table 6).
- Nationally, fewer adults have received a COVID-19 vaccine in the last 12 months (9.3%; Table 6), compared to the 12 months prior (9.8% from 15 April 2024 to 13 April 2025).
- In the last 12 months, vaccine coverage decreased in all age groups, with the largest decrease seen in 65–74 years age group (from 23.1% in the 12 months prior to 22.1% in the last 12 months).
- There has been substantial variation in COVID-19 vaccine coverage across age groups, ranging from 3.9% in adults aged 18–64 years to 35.8% in adults aged 75 years and over. Vaccine coverage increases with increasing age (Table 6).
- There has been some variation in vaccine coverage across jurisdictions, ranging from 3.8% in the NT to 16.2% in the ACT (Table 6).

Table 6: COVID-19 vaccine coverage*†‡ by age group and jurisdiction, Australia, 14 April 2025 to 19 April 2026

Age group	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total
Last 12 months (14 April 2025 to 19 April 2026)									
18–64 years	8.9	3.3	1.9	3.7	3.9	7.0	4.1	3.8	3.9
65–74 years	42.0	20.5	13.4	20.9	23.1	33.1	22.4	22.1	22.1
≥ 75 years	59.1	34.1	23.9	34.5	36.6	49.2	35.1	36.3	35.8
All ages (18 years and over)	16.2	8.6	3.8	8.8	10.4	16.1	9.2	8.9	9.3
Last 6 months (20 October 2025 to 19 April 2026)									
18–64 years	2.7	0.9	0.6	1.1	1.2	1.8	1.1	0.6	1.0
65–74 years	19.0	7.1	4.8	8.0	10.1	12.2	8.0	6.0	7.9
≥ 75 years	33.4	15.5	9.2	16.4	19.8	23.1	16.4	13.6	16.6
All ages (18 years and over)	6.7	3.2	1.3	3.5	4.6	6.0	3.4	2.4	3.4

Source: Australian Immunisation Register (AIR) as at 20 April 2026

* COVID-19 vaccine coverage among the general population uses the most recently available Australian Bureau of Statistics Estimated Resident Population (ERP) as denominator for population data. Age in years is calculated as at the reporting week.

† COVID-19 vaccine coverage is influenced by changes in COVID-19 vaccine recommendations and eligibility criteria. For this reason, coverage rates in the current 12 month period and previous 12 month periods may not be directly comparable. Coverage data in these tables may differ slightly from coverage estimates in other reports due to differences in calculation methodologies and/or different data download dates.

‡ Jurisdiction is based on the state or territory in which a vaccine was administered and may differ from a person's residential address. Population denominator data used to calculate COVID-19 vaccine coverage are based on an individual's residential address. Total rows will include individuals where jurisdiction was missing.

- Nationally, 1.3% of Aboriginal and Torres Strait Islander adults (aged 18 years or over) have received a COVID-19 vaccine in the last six months (Table 7).
- Nationally, fewer Aboriginal and Torres Strait Islander adults have received a COVID-19 vaccine in the last 12 months (4%; Table 7), compared to the 12 months prior (4.5% from 15 April 2024 to 13 April 2025).
- In the last 12 months, vaccine coverage decreased in all age groups of Aboriginal and Torres Strait Islander people, with the largest decrease seen in 65–74 years age group (from 16.4% in the 12 months prior to 14.6% in the last 12 months).
- Among Aboriginal and Torres Strait Islander people there has been substantial variation in COVID-19 vaccine coverage across age groups, ranging from 2.3% in adults aged 18–64 years to 23.5% in adults aged 75 years and over. Vaccine coverage increases with increasing age (Table 7).
- Among Aboriginal and Torres Strait Islander people, there has been slight variation in vaccine coverage across jurisdictions, ranging from 2.5% in the NT to 8.2% in the ACT (Table 7).

Table 7: COVID-19 vaccine coverage*†‡ among Aboriginal and Torres Strait Islander populations by age group and jurisdiction, Australia, 14 April 2025 to 19 April 2026

Age group	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total
Last 12 months (14 April 2025 to 19 April 2026)									
18–64 years	5.5	2.3	1.8	2.1	2.3	4.4	3.3	1.9	2.3
65–74 years	28.9	15.4	8.5	13.7	14.6	27.7	16.9	13.0	14.6
≥ 75 years	42.9	25.3	13.6	21.8	25.0	36.5	27.8	23.2	23.5
All ages (18 years and over)	8.2	4.3	2.5	3.6	4.1	8.1	5.5	3.2	4.0
Last 6 months (20 October 2025 to 19 April 2026)									
18–64 years	1.4	0.6	0.6	0.5	0.7	1.2	1.0	0.3	0.6
65–74 years	12.9	5.3	3.0	4.7	5.6	10.2	6.6	3.7	5.1
≥ 75 years	20.9	11.3	5.7	9.6	12.4	15.6	12.2	7.5	10.2
All ages (18 years and over)	2.7	1.5	0.9	1.1	1.5	2.7	1.9	0.7	1.3

Source: Australian Immunisation Register (AIR) as at 20 April 2026

* COVID-19 vaccine coverage among Aboriginal and Torres Strait Islander populations is based on the AIR population as known at the reporting week. Age in years is calculated as at the reporting week.

† COVID-19 vaccine coverage in the most recent 12 month period may not be directly comparable to previous 12 month periods due to changes in COVID-19 vaccine eligibility criteria. Coverage data in these tables may differ slightly from coverage estimates in other reports due to differences in calculation methodologies and/or different data download dates.

‡ Jurisdiction is based on the state or territory in which a vaccine was administered and may differ from a person's residential address. Population denominator data used to calculate COVID-19 vaccine coverage are based on an individual's residential address. Total rows will include individuals where jurisdiction was missing.

- *Influenza vaccine coverage data are not yet available for the seasonal influenza campaign in 2026. We expect to publish influenza vaccine coverage data in May 2026.*

- Since the commencement of the National RSV Mother and Infant Protection Program on 3 February 2025, 236,884 Abrysvo doses have been administered to pregnant people nationally (Table 8).
- While high maternal vaccine uptake is a positive indicator of maternal program success, it may result in lower nirsevimab uptake rates in infants. This is because maternal antibodies passed to the infant can provide protection against RSV, potentially reducing the need for infant immunisation.

Table 8: Number of doses of Abrysvo administered to pregnant people by jurisdiction*, Australia, 3 February to 19 April 2026

	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total
Age group									
15–24 years	307	5,929	531	5,668	1,302	634	3,489	2,268	20,130
25–39 years	5,417	62,788	2,108	38,211	14,202	4,445	56,232	20,664	204,069
40–54 years	360	4,126	102	1,991	808	198	3,885	1,215	12,685
Total (15–54 years)	6,084	72,843	2,741	45,870	16,312	5,277	63,606	24,147	236,884

Source: Australian Immunisation Register (AIR) as at 20 April 2026

* Jurisdiction is based on the state or territory in which a vaccine was administered and may differ from a person's residential address. Total rows will include individuals where jurisdiction was missing.

- In the last six months, 4% of infants (aged < 8 months) have received nirsevimab (Table 9).
- There has been some variation in nirsevimab uptake in infants across jurisdictions, ranging from 0.5% in SA to 11.1% in the NT (Table 9).
- The current trend is likely due to variation in the seasonality and eligibility criteria between state and territory programs, as well as the presence of previous nirsevimab programs. Some state and territory programs are seasonal (from 1 April to 30 September), whereas others are year-round. In states with seasonal programs (SA, Tas, Vic, and parts of WA), uptake may appear disproportionately lower at this time of the year.

Table 9: Nirsevimab (Beyfortus) uptake in the last six months*†‡ by age group and jurisdiction, Australia, 20 October 2025 to 19 April 2026

	ACT	NSW	NT	Qld	SA	Tas	Vic	WA	Total
Age group									
Infants (aged < 8 months)	1.4	4.2	11.1	8.9	0.5	1.5	1.2	3.8	4.0
Young children (aged ≥ 8 to 24 months)	0.1	0.1	0.2	0.0	0.1	0.2	0.1	0.1	0.1

Source: Australian Immunisation Register (AIR) as at 20 April 2026

* Reporting of RSV monoclonal antibodies to the AIR is not compulsory; therefore, uptake is likely to be underestimated. Uptake data in these tables may differ slightly from estimates in other reports due to differences in calculation methodologies and/or different data download dates.

† For infants and young children vaccinated, age in months is calculate as months between the immunisation encounter and date of birth rounded down as at the reporting date. For the infant and young children population, age in months is calculated as months between the AIR data extract date and date of birth rounded down as at the reporting date.

‡ Jurisdiction is based on the state or territory in which a vaccine was administered and may differ from a person's residential address. Total rows will include individuals where jurisdiction was missing. Population denominator data used to calculate nirsevimab uptake are based on an individual's residential address as recorded on Medicare.

Vaccine effectiveness

- Vaccine effectiveness (VE) is the reduction in risk of influenza and its complications in those vaccinated, compared to those not vaccinated.
- Interim Australian data as part of the Global Influenza Vaccine Effectiveness (GIVE) Collaboration indicate that in 2025, people who received the influenza vaccine were about 53% less likely to visit general practice or be hospitalised with influenza compared to those who were unvaccinated. Please note, these interim estimates were based on incomplete data, and the final VE estimates - expected to be released in the 2025 Annual Australian Respiratory Surveillance Report later in 2026 - may change.
- It is too early to assess VE for the 2026 influenza season.

Vaccine match

- In the year to date, 100% (32/32) of influenza A(H1N1) isolates, 87.9% (240/273) of influenza A(H3N2) isolates and 88.9% (32/36) of influenza B/Victoria lineage isolates characterised have been antigenically similar to the corresponding 2026 southern hemisphere vaccine components.

2026 southern hemisphere vaccine composition

The composition of influenza vaccines for Australia in 2026 differs from the 2025 southern hemisphere and 2025/26 northern hemisphere composition. The southern hemisphere 2026 vaccine contains 2 new strains for the influenza A(H1N1)pdm09 and A(H3N2) subtype virus components.

The following influenza viruses are used for the 2026 southern hemisphere trivalent influenza vaccines in Australia:

Egg-based influenza vaccines:

- an A/Missouri/11/2025 (H1N1)pdm09-like virus
- an A/Singapore/GP20238/2024 (H3N2)-like virus
- a B/Austria/1359417/2021 (B/Victoria lineage)-like virus. /

Cell-based influenza vaccines:

- an A/Missouri/11/2025 (H1N1)pdm09-like virus
- an A/Sydney/1359/2024 (H3N2)-like virus
- a B/Austria/1359417/2021 (B/Victoria lineage)-like virus.

The continued absence of confirmed detection of naturally occurring B/Yamagata lineage viruses after March 2020 is indicative of a very low risk of infection by B/Yamagata lineage viruses. Since September 2023, the WHO has recommended that the inclusion of a B/Yamagata lineage antigen in seasonal influenza vaccines is no longer warranted.